

COMMON FIXED POINT THEOREM OF AN INFINITE SEQUENCE OF MAPPINGS IN HILBERT SPACE

ARVIND KUMAR SHARMA¹, V. H. BADSHAH² & V. K. GUPTA³

¹Department of Engineering Mathematics, M.I.T.M, Ujjain, Madhya Pradesh, India

²School of Studies in Mathematics, Vikram University, Ujjain, Madhya Pradesh, India

³Department of Mathematics, Government Madhav Science College, Ujjain, Madhya Pradesh, India

ABSTRACT

The aim of this present paper is to obtain a common fixed point for an infinite sequence of mappings on Hilbert space. Our purpose here is to generalize the our previous result [7]

KEYWORDS: Common Fixed Point, Hilbert Space, Infinite Sequence of Mappings

2010 Mathematics Subject Classification: 47H10.

1. INTRODUCTION

In 2011, Sharma Badshah and Gupta [7] have proved common fixed point theorem for the sequence $\{T_n\}_{n=1}^{\infty}$ of mappings satisfying the condition

$$\|T_i x - T_j y\| \leq \alpha \frac{\|x - T_i x\|^2 + \|y - T_j y\|^2}{\|x - T_i x\| + \|y - T_j y\|} + \beta \|x - y\| \quad (A)$$

for all $x, y \in S$ and $x \neq y$; $\alpha \geq 0$, $\beta \geq 0$ and $2\alpha + \beta < 1$.

In 2005, Badshah and Meena [1] have proved common fixed point theorem for the sequence $\{T_n\}_{n=1}^{\infty}$ of mappings satisfying the condition

$$\|T_i x - T_j y\| \leq \alpha \frac{\|x - T_i x\| \cdot \|y - T_j y\|}{\|x - y\|} + \beta \|x - y\| \quad (B)$$

for all $x, y \in S$ with $x \neq y$ also $\alpha \geq 0$, $\beta \geq 0$ and $\alpha + \beta < 1$.

In 1991, Koparde and Waghmode [3] have proved common fixed point theorem for the sequence $\{T_n\}_{n=1}^{\infty}$ of mappings satisfying the condition

$$\|T_i x - T_j y\|^2 \leq a \left(\|x - T_i x\|^2 + \|y - T_j y\|^2 \right) \quad (C)$$

for all $x, y \in S$ and $x \neq y$; $0 \leq a < 1/2$

Later in 1998, Pandhare and Waghmode [5] have proved common fixed point theorem for the sequence $\{T_n\}_{n=1}^{\infty}$ of mappings satisfying the condition

$$\|T_i x - T_j y\|^2 \leq a \|x - T_i x\|^2 + b \left(\|x - T_i x\|^2 + \|y - T_j y\|^2 \right) \quad (D)$$

for all $x, y \in S$ and $x \neq y$; $0 \leq a, 0 \leq b < 1$ and $a + 2b < 1$.

This result is generalizes by Veerapandi and Kumar [7] and the new condition is

$$\|T_i x - T_j y\|^2 \leq a \|x - y\|^2 + b \left(\|x - T_i x\|^2 + \|y - T_j y\|^2 \right) + \frac{c}{2} \left(\|x - T_i x\|^2 + \|y - T_j y\|^2 \right) \quad (E)$$

for all $x, y \in S$ and $x \neq y$ where $0 \leq a, b, c < 1$ and $a + 2b + 2c < 1$.

Now we introduce a new condition for the generalization of following known results.

Theorem 1: [7] Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be an infinite sequence of mappings satisfy (A). Then $\{T_n\}_{n=1}^{\infty}$ has a unique common fixed point.

Theorem 2: [1] Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be an infinite sequence of mappings satisfy (B). Then $\{T_n\}_{n=1}^{\infty}$ has a unique common fixed point.

Theorem 3: [3] Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be a sequence of mappings satisfy (C). Then $\{T_n\}_{n=1}^{\infty}$ has a unique common fixed point.

Theorem 4: [5] Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be a sequence of mappings satisfy (D). Then $\{T_n\}_{n=1}^{\infty}$ has a unique common fixed point.

Theorem 5: [8] Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be a sequence of mappings satisfy (E). Then $\{T_n\}_{n=1}^{\infty}$ has a unique common fixed point.

Main Result

We proved fixed point theorem for the infinite sequence $\{T_n\}_{n=1}^{\infty}$ to generalize our previous results [7].

Theorem: Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be an infinite sequence of mappings satisfying the following condition

$$\|T_i x - T_j y\| \leq \left(\alpha + \beta \frac{\|x - T_i x\|}{\|x - y\|} \right) \|y - T_j y\| \quad (F)$$

for all $x, y \in S$ and $x \neq y$; $\alpha \geq 0, \beta \geq 0$ and $2\alpha + \beta < 1$.

Then $\{T_n\}_{n=1}^{\infty}$ has a unique common fixed point.

Proof: Let S be a closed subset of a Hilbert space H and $\{T_n\}_{n=1}^{\infty} : S \rightarrow S$ be an infinite sequence of mappings.

Let $x_0 \in S$ be any arbitrary point in S .

Define a sequence $\{x_n\}_{n=1}^{\infty}$ in S by

$$x_{n+1} = T_{n+1}x_n, \text{ for } n = 0, 1, 2, \dots$$

For any integer $n \geq 1$

$$\begin{aligned} \|x_{n+1} - x_n\| &= \|T_{n+1}x_n - T_nx_{n-1}\| \\ &\leq \left(\alpha + \beta \frac{\|x_n - T_{n+1}x_n\|}{\|x_n - x_{n-1}\|} \right) \|x_{n-1} - T_nx_{n-1}\| \\ &\leq \left(\alpha + \beta \frac{\|x_n - x_{n+1}\|}{\|x_n - x_{n-1}\|} \right) \|x_{n-1} - x_n\| \\ &\leq \alpha \|x_{n-1} - x_n\| + \beta \|x_n - x_{n+1}\| \\ \text{i.e. } \|x_{n+1} - x_n\| &\leq \alpha \|x_{n-1} - x_n\| + \beta \|x_n - x_{n+1}\| \\ \Rightarrow (1 - \beta) \|x_{n+1} - x_n\| &\leq \alpha \|x_n - x_{n-1}\| \\ \Rightarrow \|x_{n+1} - x_n\| &\leq \frac{\alpha}{1 - \beta} \|x_n - x_{n-1}\| \end{aligned}$$

If $k = \frac{\alpha}{1 - \beta}$ then $k < 1$.

$$\begin{aligned} \|x_{n+1} - x_n\| &\leq k \|x_n - x_{n-1}\| \\ &\leq k \|x_n - x_{n-1}\| \leq k^2 \|x_{n-1} - x_{n-2}\| \leq k^3 \|x_{n-2} - x_{n-3}\| \leq \dots \leq k^n \|x_1 - x_0\| \\ \text{i.e. } \|x_{n+1} - x_n\| &\leq k^n \|x_1 - x_0\| \text{ for all } n \geq 1 \text{ is integer.} \end{aligned}$$

Now for any positive integer $m \geq n \geq 1$

$$\begin{aligned} \|x_n - x_m\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - x_{n+2}\| + \dots + \|x_{m-1} - x_m\| \\ &\leq k^n \|x_1 - x_0\| + k^{n+1} \|x_1 - x_0\| + \dots + k^{m-1} \|x_1 - x_0\| \\ &\leq k^n \|x_1 - x_0\| (1 + k + \dots + k^{m-n-1}) \\ \text{i.e. } \|x_n - x_m\| &\leq \left(\frac{k^n}{1 - k} \right) \|x_1 - x_0\| \rightarrow 0 \text{ as } n \rightarrow \infty \text{ (} k < 1 \text{)} \end{aligned}$$

Therefore $\{x_n\}_{n=1}^{\infty}$ is a Cauchy sequence.

Since S is a closed subset of a Hilbert space H , so $\{x_n\}_{n=1}^{\infty}$ converges to a point u in S .

Now we will show that u is common fixed point of infinite sequence $\{T_n\}_{n=1}^{\infty}$ of mappings from S into S .

Suppose that $T_n u \neq u$ for all n .

Consider for any positive integer m ($\neq n$)

$$\begin{aligned}
 \|u - T_m u\| &\leq \|u - x_n\| + \|x_n - T_m u\| \\
 &= \|x_n - T_m u\| \\
 &= \|T_n x_{n-1} - T_m u\| \\
 &\leq \left(\alpha + \beta \frac{\|x_{n-1} - T_n x_{n-1}\|}{\|x_{n-1} - u\|} \right) \|u - T_m u\| \\
 &\leq \left(\alpha + \beta \frac{\|x_{n-1} - x_n\|}{\|x_{n-1} - u\|} \right) \|u - T_m u\| \\
 &\leq \alpha \|u - T_m u\| + \beta \frac{\|x_{n-1} - x_n\|}{\|x_{n-1} - u\|} \|u - T_m u\| \\
 \text{i.e.} \quad \|u - T_m u\| &\leq \alpha \|u - T_m u\| + \beta \frac{\|x_{n-1} - x_n\|}{\|x_{n-1} - u\|} \|u - T_m u\| \\
 \Rightarrow \|u - T_m u\| &\leq \frac{\beta}{1 - \alpha} \frac{\|x_{n-1} - x_n\|}{\|x_{n-1} - u\|} \|u - T_m u\| \rightarrow 0 \text{ as } n \rightarrow \infty
 \end{aligned}$$

$$\text{So } \|u - T_m u\| \leq 0.$$

Hence $u = T_m u$ and so $u = T_n u$ for all n .

Hence u is a common fixed point of infinite sequence $\{T_n\}_{n=1}^{\infty}$ of mappings.

Uniqueness

.Suppose that there is $u \neq v$ such that $T_n v = v$ for all n .

$$\text{Consider } \|u - v\| = \|T_n u - T_n v\|$$

$$\leq \left(\alpha + \beta \frac{\|u - T_n u\|}{\|u - v\|} \right) \|v - T_n v\|$$

$$\text{i.e. } \|u - v\| \leq 0$$

$$\Rightarrow \|u - v\| = 0$$

Thus $u = v$.

Hence fixed point is unique.

Example: Let $X = [0, 1]$, with Euclidean metric d . Then $\{X, d\}$ is a Hilbert space with the norm defined by $d(x, y) = \|x - y\|$

Let $\{x_n\}_{n=1}^{\infty} = \left\{\frac{1}{2^n}\right\}_{n=1}^{\infty}$ be the sequence in X and let $\{T_n\}_{n=1}^{\infty}$ be the infinite sequence of mappings such that

$$x_{n+1} = T_{n+1}x_n, \text{ for } n = 0, 1, 2, \dots$$

Taking $x = \frac{1}{2^n}$ and $y = \frac{1}{2^{n-1}}$; $x \neq y$. Also $i = n+1$ and $j = n$.

Then from (F) $\{x_n\}_{n=1}^{\infty}$ is a Cauchy sequence in X , which is converges in X also it has a common point in X .

CONCLUSIONS

The theorem proved in this paper by using rational inequality is improved and stronger form of some earlier inequality given by Badshah and Meena [1], Sharma Badshah and Gupta [7], Koparde and Waghmode [3], Pandhare and Waghmode [5], Veerapandi and Kumar [8].

REFERENCES

1. Badshah, V.H. and Meena, G., (2005): Common fixed point theorems of an infinite sequence of mappings, Chh. J. Sci. Tech. Vol. 2, 87-90.
2. Browder, Felix E. (1965): Fixed point theorem for non compact mappings in Hilbert space, Proc Natl Acad Sci U S A. 1965 June; 53(6): 1272-1276.
3. Koparde, P.V. and Waghmode, B.B. (1991): On sequence of mappings in Hilbert space, The Mathematics Education, XXV, 197.
4. Koparde, P.V. and Waghmode, B.B. (1991): Kannan type mappings in Hilbert spaces, Scientist Phyl. Sciences Vol. 3, No. 1, 45-50.
5. Pandhare, D.M. and Waghmode, B.B. (1998): On sequence of mappings in Hilbert space, The Mathematics Education, XXXII, 61.
6. Sangar, V.M. and Waghmode, B.B. (1991): Fixed point theorem for commuting mappings in Hilbert space-I, Scientist Phyl. Sciences Vol. 3, No.1, 64-66.
7. Sharma, A.K., Badshah, V.H and Gupta, V.K. (2012): Common fixed point theorems of a sequence of mappings in Hilbert space, Ultra Scientist Phyl. Sciences.
8. Veerapandhi, T. and Kumar, Anil S. (1999): Common fixed point theorems of a sequence of mappings in Hilbert space, Bull. Cal. Math. Soc. 91 (4), 299-308.

